



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
24.05.2000 Bulletin 2000/21

(51) Int. Cl.⁷: **C12N 15/82, C12N 15/87**

(21) Application number: **91301545.9**

(22) Date of filing: **26.02.1991**

(54) **Plant transformation process with early identification of germ line transformation events**

Pflanzentransformationsverfahren mit Früherkennung von Keimbahn-Transformationsereignissen

Procédé de transformation de plantes avec détection précoce des événements de transformation de la lignée germinale

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(30) Priority: **26.02.1990 US 485111**

(43) Date of publication of application:
04.09.1991 Bulletin 1991/36

(73) Proprietor: **MONSANTO COMPANY**
St. Louis, Missouri 63167 (US)

(72) Inventors:
• **Christou, Paul**
Madison, Wisconsin 53717 (US)
• **McCabe, Dennis E.**
Middleton, Wisconsin 53562 (US)

(74) Representative:
Bentham, Stephen et al
J.A. KEMP & CO.
14 South Square
Gray's Inn
London WC1R 5LX (GB)

(56) References cited:
EP-A- 0 301 749

- **ABSTRACTS VIITH INTERNATIONAL CONGRESS ON PLANT TISSUE AND CELL CULTURE June 1990, AMSTERDAM, THE NETHERLANDS, page 51; CHRISTOU, P., ET AL.: 'Commercialization of a process for the introduction of foreign genes in elite cultivars of soybean using electric discharge acceleration'**
- **PLANT MOLECULAR BIOLOGY. vol. 14, no. 2, February 1990, DORDRECHT, THE NETHERLANDS. pages 261 - 268; TOMES, D. T., ET AL.: 'Transgenic tobacco plants and their progeny derived by microprojectile bombardment of tobacco leaves'**
- **PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA. vol. 86, no. 19, October 1989, WASHINGTON US, pages 7500 - 7504; CHRISTOU, P., ET AL.: Inheritance and expression of foreign genes in transgenic soybean plants'**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention relates to the genetic transformation of bean plants in general, and relates, in particular, to the efficient germ line transformation of soybean plants utilizing particle-mediated techniques for plant transformation.

[0002] It is now becoming possible to introduce into some of the major crop plants foreign genes of potential interest in those plants. This process of genetic engineering has been accomplished with certain model species, such as tobacco, petunia, and carrot, and has now been accomplished in such major crops species soybean and cotton. In procedures for the genetic engineering of plants, it is desired that the genetic transformation of the plant tissues be of the germ line of the plant tissues. Germ line refers to the inheritable genetic material of the plant which is permanently altered by the transformation process in such a fashion that the plant will pass to its progeny by normal genetic inheritance the inserted foreign gene. While the transformation of somatic, or non-germ line cells, may be desired in some instances, in general in the genetic engineering of crop plants it is desired that germ line transformation of plant lines be achieved as quickly and efficiently as possible.

[0003] The most common previously utilized technique for the genetic engineering of plants involves the use of the soil-dwelling plant pathogenic bacterium Agrobacterium tumefaciens. A. tumefaciens has the natural ability to transfer a portion of its DNA, referred to as T-DNA, into the genome of susceptible plant cells. By changing the native T-DNA in an Agrobacterium strain, it is possible to use this unique trait of Agrobacterium to transfer desired genes into single plant cells. If the introduced gene includes a selectable marker, such as a herbicide or antibiotic resistance trait, it is possible thereafter to select for transformed cells in a tissue culture by imposing the putative transformant tissues to the selection pressure of the appropriate antibiotic or herbicide. Unfortunately, in soybean almost all cultivars are resistant to Agrobacterium infection and are thus very resistant to transformation with Agrobacterium. In addition, antibiotic resistant markers, such as kanamycin resistance commonly used in Agrobacterium plant transformation procedures with other plant species, have been found to be of limited utility in soybean transformation experiments. Accordingly, while it may be possible to utilize Agrobacterium-mediated transformation techniques in soybean, it is a difficult endeavor because of the lack of effective selectable markers.

[0004] Other techniques for transforming plants do exist, however. In particular, there exists a general approach to the transformation of plant cells which is based on delivering the transforming DNA into the plant cells by coating the DNA onto small inert carrier particles which are physically hurled into the target plant tissues. The technique of particle-mediated plant cell transformation was first demonstrated with somatic cells in such tissues as the epidermal tissue of onion and other such model cell cultures. Klein et al., Nature, 327:70-73 (1987). Later the originators of the particle-mediated transformation technique were able to achieve genetically engineered tobacco plants by the transformation of tobacco in tissue culture, using a selectable marker, which was then subsequently regenerated into whole plants. Klein et al., Proc. Natl. Acad. Sci. USA, 85:8502-8505 (1988). Rather than attempting to use particle-mediated transformation techniques on plant cells in culture, another approach was developed in which the growing meristems of plants were subjected to a particle-mediated transformation event. From such a technique, stable transformation of the germ line of soybean plants was achieved. McCabe et al., Bio/Technology, 6:923-926 (1988) see also Christou et al., Proc. Natl. Acad. Sci., 1989, vol 86, pp 7500-7504. This technique is not dependent on the availability of a selectable marker for the plant species.

[0005] In developing the technique for the germ line transformation of soybean plants using a particle-mediated technique based on meristem transformation, it was discovered that the transformation events often resulted in chimeric plants, which are plants in which some, but not all, of the tissues had been genetically transformed with the introduced DNA. McCabe et al., supra. Although the technique was thus useful to create genetically engineered plants, it was somewhat burdensome in the sense that large numbers of tissues had to be subject to transformation events, and large numbers of plants had to be cultivated from the putative transformed tissues in order to discover those particular shoots and plants which properly expressed the introduced DNA. Therefore, the ability to identify transformation events early in the process giving rise to heritable germ line transformation of tissues creates the ability to effectuate dramatic savings in the practical and cost-effective genetic transformation of plants resulting in reduced labor costs and reduced time and energy expended in cultivating the non-transformed tissues which were subject to the transformation events.

[0006] In seeking a germ line transformation of a plant species, such as soybean, it would be helpful if the progenitor tissue of the germ cells of the plant were identified in the growing meristem or shoot. Unfortunately, the science of developmental morphology of plant cells has not developed to the point that the ancestor cells of the soybean germ cells is known. Accordingly, if it is a plant meristem or embryo that is being transformed, no present knowledge exists as to which precise cells in that meristem or embryo must be transformed to achieve germ line transformation. Therefore, any correlation between categories of cells transformed in a growing soybean plant and a germ line transformation event would have to be determined empirically.

[0007] The present invention provides a method of creating germ line transformed soybean plants comprising the steps of:

(a) preparing an exogenous genetic construction suitable for insertion into soybean plants, which construction comprises expression constructs for both a marker gene and a gene of interest, each of which comprise a coding sequence and a flanking regulatory sequence effective in soybean cells to produce a gene product coded by the coding sequence in the cells of soybean plants;

- 5 (b) coating copies of the exogenous genetic construction onto carrier particles of biologically inert material, which are small in size relative to the size of soybean cells;
- (c) accelerating the coated carrier particles into the cells of regenerable soybean tissue;
- (d) regenerating shoots from the soybean tissue;
- (e) severing a segment of the stem from each of the shoots;
- 10 (f) assaying the stem segments for the expression of the gene product of the marker gene;
- (g) regenerating into whole sexually mature soybean plants only the shoots, the stem segments of which expressed the gene product of the marker gene in at least a major portion of its pith;
- (h) sexually propagating the whole plants to create progeny plants; and
- 15 (i) assaying for the expression of the gene product of the marker gene in the progeny plants to indicate the germ line insertion of the gene of interest into the germ line of the progeny plants.

[0008] The invention further provides a method for the germ line transformation of bean plants, such as soybean plants, which method comprises the steps of:

- 20 (a) preparing an exogenous genetic construction comprising a coding sequence and a flanking regulatory sequence effective to express a gene product coded by the coding sequence in the cells of the plant;
- (b) coating copies of the exogenous genetic construction onto carrier particles which are of a biologically inert material and which are small in size relative to the size of the cells of the plant;
- (c) accelerating the coated carrier particles into the cells of a regenerable tissue of the plant;
- 25 (d) regenerating the tissue into plants;
- (e) during the regeneration process, assaying a selected portion of the stem of each regenerating plant for the presence of the genetic construction, the selected stem portion including a pith section, the assay determining whether the genetic construction is expressed in the pith section;
- (f) sexually multiplying preferentially the plants the selected portions of which assay positive for the presence of the
- 30 (g) verifying the presence of the genetic construction in the progeny plants.

[0009] It is an object of the present invention to provide a method of particle-mediated transformation of plants which is inherently more efficient and cost-effective than previously available techniques.

35 **[0010]** It is an object of the present invention to make more cost effective the genetic engineering of plants, and soybean plants in particular, by identifying, at the earliest stage possible in the process, those plants which are most likely to yield germ line transformants so that only those plants need to be cultivated into mature plants.

[0011] It is another object of the present invention to optimize the use of reporter or marker genes in plant transformation experiments even in the absence of selectable markers to make possible the genetic engineering of plants in

40 those plants for which reliable dominant selectable markers are not available.

[0012] Other objects, advantages, and features of the present invention will become apparent from the following specification when taken in conjunction with the accompanying drawings.

45 Fig. 1 is a schematic illustration of an apparatus suitable for use in the particle-mediated transformation of plant tissues.

Fig. 2 is a top plan view of the spark discharge chamber of the apparatus of Fig. 1.

Fig. 3 is a highly stylized illustration of a cross-section through a soybean stem.

Figs. 4 - 8 are other stylized illustrations of a cross-section through a soybean stem showing various phenotype of marker gene expression.

50 **[0013]** In accordance with the method of the present invention, a series of indices are described which are used to identify high likelihood germ line transformation events among the shoots and plants produced from a particle-mediated plant transformation process. These indices are based on a classification of certain phenotypes of expression of a marker gene in tissues of a regenerating plant at certain defined stages.

55 **[0014]** To fully understand the advantages of the present method, it is helpful to appreciate certain considerations about the nature of accelerated particle-mediated transformation of plants. Since the transforming DNA is carried into the plant tissues on particles, and since the number of particles must be limited to avoid destroying the plant tissues transformed, only a percentage of the cells in a treated plant tissue will receive transforming DNA, and only a percent-

age of those cells will be transformed. Therefore, in the absence of a selection agent to kill non-transformed cells preferentially, the result of regenerating plants from such treated tissues will be plants most of the tissues of which are not transformed. It has been found that even of the plants which are at least in part transformed, most of the transformations do not result in germ line transformations. This invention is therefore directed to identifying those desired germ line transformation events at an early stage so that regeneration of the non-germ line plants can be minimized.

[0015] Thus it is not necessary to use the early germ line identification process described here to achieve a germ line transformation of soybean. It is possible to regenerate all plants recovered from the treated tissue, sexually propagate all the plants and assay all the progeny. The drawback to this approach is that most of the effort in the regeneration and propagation process will be wasted on non-germ line transformation events. The present invention helps to avoid that waste and thereby assist in the efficient creation of lines of genetically transformed soybean.

[0016] The present invention is thus based on a technique involving the particle-mediated transformation of plant cells. Therefore, to better understand the context of the present invention, it is necessary to understand the general technique of particle-mediated transformation of plant cells and the apparatus which may be used therefore.

[0017] In the process of particle-mediated transformation of plant cells, a carrier particle consisting of a small inert relatively dense particle of material is coated with a transforming genetic construction of DNA and then is physically accelerated so that it is delivered into the interior of the growing meristematic or embryonic cells to be transformed. The transforming DNA is thus carried into individual cells, but the carrier particles are small enough such that the individual cells are neither destroyed nor seriously incapacitated. It has been found that by delivering DNA on such carrier particles in such a fashion into the cells of plants, such as soybeans, whole transformed germ line plants can be obtained to result in transgenic plants in plant lines.

[0018] There are several factors which are necessary to be considered in the creation of germ line plant transformations in these fashions. The genetic construction must be one properly constructed to be expressed in plant tissues. The apparatus utilized must be of a type capable of delivering the carrier particles with the coated DNA on them into plant cells in such a fashion that a suitable number of cells are transformed. There are many types of mechanical systems which can be envisioned to accelerate biologically inert small carrier particles. Possible mechanisms include ballistic explosive acceleration of particles, centrifugal acceleration of particles, electrostatic acceleration of particles, or any other analogous system capable of providing momentum and velocity to small inert particles. The mechanism used herein to achieve particle-mediated plant transformation is based on an adjustable electric voltage spark discharge device. The apparatus is illustrated in a schematic fashion in Figs. 1 and 2.

[0019] The particle-acceleration apparatus is generally indicated at 10 in Fig. 1. The apparatus consists of a spark discharge chamber 12 into which are inserted two electrodes 14 which are spaced apart by a distance of approximately one to two millimeters. The spark discharge chamber 12 is a horizontally extending rectangle having two openings 16 and 18 extending out its upward end. The opening 16 is covered by an access plate 20. The opening 18, located on the side of the rectangle of the spark discharge chamber 12 opposite from the electrodes 14, is intended to be covered by a carrier sheet 22. The electrodes 14 are connected to a suitable adjustable source of electric discharge voltage. Such a source of electric discharge voltage would preferably include suitable electric switching connected to a capacitor of the one to two microfarad size range, with the amount of the voltage of the charge introduced onto the capacitor being adjustable, such as through the use of an autotransformer, through a range of perhaps 1 to 50,000 volts. Suitable high voltage electric switching (not shown) is provided so that the capacitor can safely be discharged through the electrodes 14 so that the apparatus can be used conveniently by a user.

[0020] The carrier sheet 22 intended to be placed upon the opening 18 in the spark discharge chamber 12 is a planar sheet of relatively stiff material, such as a sheet of aluminized saran coated mylar. Above the opening 18 in the discharge chamber 12, positioned approximately 15 millimeters above it, is a retaining screen 24. Placed above the retaining screen 24 at a distance of approximately 5 to 25 millimeters above the retaining screen, is a target surface 26. The target surface 26 can be any suitable culture surface onto which the material to be transformed may readily be placed such as, most conveniently, an overturned petri dish into which the plant tissues have been positioned for culture. Copies of the exogenous foreign genetic construction intended to be transformed into the plant tissues are prepared by suitable DNA preparation techniques well known to those of ordinary skill in the art, and multiple copies of the genetic construction are made. The copies of the foreign genetic construction, in aqueous solution, are then coated onto small particles of a durable dense biologically inert carrier material, such as gold, the carrier particles typically being in a size range of 1 to 3 microns. The carrier particles with the exogenous genetic construction thereon are then placed upon the carrier sheet 22, which is inserted at the proper opening on the top of the spark discharge chamber 12. The target surface 26, including the living plant material thereon, is then placed in position above the retaining screen 24. A small droplet of water, 30, preferably about 10 microliters in size, is then placed bridging between the ends of the electrodes 14. The access cover 20 is placed in position on top of the spark discharge chamber 12. At this point the entire apparatus is enclosed in a vacuum chamber and a vacuum is drawn until it is in the range of approximately 500 millimeters of mercury. As the vacuum is being drawn, a supply of helium is bled into the vacuum chamber, replacing the remaining atmosphere in the chamber with helium. The lower relative density of helium, combined with the

reduced pressure, helps to lower the drag on the gold particles.

[0021] At this point, the spark discharge between the electrodes 14 may be initiated by the user. This is done by means of the appropriate electric switching which applies the voltage stored in the capacitor across the terminals of the electrodes 14. The force of this electric discharge bridges the spark discharge gap between the electrodes 14 instantly vaporizing the small droplet of water previously placed therebetween. The force of the vaporization of that water creates a shockwave within the spark discharge chamber 12 which radiates outward in all directions. The impact of the radiating shockwave upon the carrier sheet 22 propels the carrier sheet 22 upward with great velocity. The upwardly traveling carrier sheet 22 accelerates until it contacts the retaining screen 24. The use of the helium within the vacuum containment for the apparatus provides less drag on the flight of the carrier sheet 22 as well as the carrier particles. At the retaining screen 24, the carrier sheet 22 is retained, and the carrier particles coated with the exogenous genetic construction previously coated thereon fly off of the carrier sheet and travel freely onward toward the target tissues. The small carrier particles then proceed into the cells of the target tissues placed on the target surface 26, and pass freely into the cytosol of the cells placed thereon. The actual momentum of the carrier particles as they impact the surface of the target tissues is adjustable, based upon the voltage of the initial electric discharge applied to the electrodes 14. Thus, by varying the amount of the electric discharge applied across the electrodes 14, the velocity by which the particles impact the target can be adjusted, and thus the depth of penetration of the carrier particles into the tissue of the target tissues can be adjusted continuously throughout the range of adjustment provided for the electric voltage applied across the electrodes 14.

[0022] To be useful in a particle-mediated transformation technique, the transforming exogenous genetic construction must be capable of performing some useful function in the cells of target plant tissues. The transforming genetic construction, which will normally be a chimeric construction in the sense that its DNA originates from more than one organism, should be capable of expressing in the target tissues gene product such as a foreign protein of interest or an antisense RNA strand. Such foreign genetic constructions, usefully embodied in expression cassette vectors for use in plant cells, are known to those of ordinary skill in the art. Typically such plant expression cassette vectors includes, besides the coding sequence of the desired exogenous or foreign gene, appropriate flanking regulatory sequences suitable for expression of the foreign gene in the plant cells, such as a promoter sequence capable of initiating transcription and a translational terminator to terminate translation of a message if protein synthesis is desired. It has been previously demonstrated that typical promoters and transcription terminators found to be effective in other plant tissues are effective in soybean as well.

[0023] The transforming genetic construction may also include a marker gene. Such a marker gene need not be a selectable marker. It need only be a marker the presence of which can be assayed using a minimal amount of plant tissue, by appropriate biochemical or phenotypic trait which may be observed in transgenic plant tissues. Such a marker gene may be the transforming DNA itself, if a completely biochemical assay for the presence of the transforming DNA is used, such as a polymerase chain reaction type of assay for the DNA itself. One convenient type of marker gene capable of detection by a phenotypic assay is the GUS gene described by Jefferson et al., Embo J., 6:3901-3907 (1987). The GUS gene, coding for the enzyme beta-glucuronidase which can be expressed in plant cells and the expression of which, in a tissue-destructive assay, will turn a convenient substraight, indigo-glucuronide, or 5-bromo-4-chloro-3-indolyl glucuronide blue in color in an in situ assay in plant tissues. Thus, the use of the GUS gene provides a convenient colorimetric assay for the expression of introduced DNA by phenotypic analysis in transformant plant tissues. Thus, in a typical transformation procedure, the desired gene of interest would be coupled in tandem in a single genetic construction, a DNA strand, with a GUS gene, and then the detection of the transforming DNA in plant tissues would be done by phenotypic analysis for the expression of the GUS enzyme in the target plant tissues.

[0024] Several plant tissues of soybean may be genetically transformed by such a particle-mediated transformation technique using the apparatus of Fig. 1. It has been found most conveniently that the excised embryonic axes from immature or mature soybean seeds may be readily transformed utilizing the procedure. The embryonic axes are excised from the soybean seeds and the primary leaves are removed to expose the meristem of the embryo. The axes are then plated on target plates containing 1% water-agar. The plates are then used as a target surface in the apparatus of Fig. 1 for a particle-mediated transformation event. The axes may then be plated in the dark on MS basal medium as modified by Barwale et al., Planta, 167:473-481 (1986) for regenerating zygotic embryos. This particular medium contains a high level of benzylaminopurine which induces multiple shoot formation in the plated embryonic tissues. Following incubation for one to two weeks in the dark, the tissues are transferred to the same basal medium with lower concentrations of benzylaminopurine, and then are cultivated in the light to promote shoot elongation. Using this technique, multiple shoots will be derived from the primary and axillary meristems on the embryonic tissues. These shoots may then be grafted onto soybean roots, or induced to form roots, to regenerate whole intact and sexually mature soybean plants. It is these shoots, and the soybean plants regenerated therefrom, which are subject to the screening techniques described herein so as to result in high numbers of germ line transformant plants.

[0025] When the soybean shoots or plantlets resulting from tissues subject to a transformation process, such as has been described above, are regenerated into whole mature plants, it has been found that only a fraction of the result-

ing soybean plants will prove to be genetically transformed with the foreign DNA. The plants which are regenerated from the shoots resulting from this tissue are referred to as R0 plants, while their progeny in successive generations are referred to as R1 and R2, etc. In addition to having a low frequency of genetic transformation within the R0 plant generation, of the R0 plants which exhibit some genetic transformation, most of the plants will be chimeric. By the use of the term chimeric in this sense, it is intended to signify that the plants will be composed of tissues which are not genetically identical, i.e., the plants will have only a portion or fraction of their tissues transformed, while the remainder of the tissues are not genetically transformed. Since it is the object of the plant transformation process such as that described herein to create stably transformed plants which carry the foreign DNA in their germ plasm, and are capable of passing the inserted DNA into their progeny, it is necessary then to implement a screening procedure to ascertain how to recover the germ line transformation events from a number, and perhaps a large number, of putatively transformed R0 plants or plant tissues.

[0026] The method disclosed herein is intended to enable the screening of R0 shoots and plants from the putatively transformed plant culture to isolate those plants which will give rise to, or at least be most likely to give rise to, progeny plants which have had their germ line transformed. In this regard it should be noted that for the progeny grown from the R0 plants, that is to say from the R1 generation onward, if the progeny plants contained the inserted foreign DNA at all, they are clonal, that is to say, non-chimeric and have their germ line cells transformed. While events are observed in which the inserted DNA appears to be present in a clonal fashion in the R0 plant, but then is found completely absent in the R1 and R2 plants, if the gene is found to be present in the R1 plant, it will prove to be stably inheritable thereafter in those of the progeny of the R1 plant which inherit the inserted gene as a trait.

[0027] The present method is a screening technique rather than a selection technique. By that it is meant that the plants, or portions of the plants, are screened for the presence of certain marker characteristics, but are not subjected to a selection criteria such as would be achieved by antibiotic or herbicide resistance routines. In part the use of such a screening technique, as opposed to a selecting technique, is required due to the absence of reliable selectable marker gene effective in soybean tissues. The preferred marker gene, as mentioned, to be the object of the screening process, is the GUS gene. The GUS assay is, however, destructive of the plant tissues and thus cannot be performed *in vivo*. It must therefore be performed on a portion of the tissues recovered from the plants to be screened and the portion of the tissue selected must be one that balances the need to gain an accurate reflection of the prospects for germ line transformation on the particular plant with the need to maintain the vigor of the plant tissues involved so that the very events sought are not lost during the assay process.

[0028] The method of the present invention is based on the selection of certain strategically located portions of plant tissue from the R0 plants during the regeneration process. In particular, three indices of the state of transformation of the R0 plants are utilized. The first index is a section of stem segment from the R0 shoot collected from the shoot prior to rooting. The second index is a section of the leaf phenotype at the first trifoliate stage of development of the seedling. The third index is a petiole/mid-rib section of a trifoliate leaf in a more mature plant, at the third or fourth trifoliate leaf stage. By doing histochemical assays for the enzyme coded by the GUS gene in each of these three stages, it is possible to predict with a high degree of confidence the germ line transformation events which have occurred. In fact, it is possible at a relatively early stage in the procedure to discard most (as many as 90-99%) of the regenerating R0 shoots and plantlets to concentrate on the remaining shoots or plantlets which have a high probability of yielding germ line transformation events. While the discarding of many such shoots or plantlets may seem wasteful, and will almost certainly involve the discarding of occasional truly transformed plants, the amount of labor and time saved by not taking large numbers of non-transformed plants to seed is large in comparison to the time involved in creating additional transformation events. Thus the laborious part of the process, i.e., taking the putatively transformed plants to seed and growing and testing the progeny, can be reduced to a minimum by the screening methodology disclosed herein.

[0029] The first index analysis of the transformed plants occurs when regeneration of shoots commences from the originally transformed plant explant. As the regenerating soybean shoots reach a finite and defined size, typically approximately 2 centimeters in size, the shoots may be isolated from the original explant and prepared for propagation, either by grafting or by hormone treatment to induce root formation. However, at the time the shoots are separated from the original explant, it is convenient at that point to take a relatively small segment of the stem from the basal portion of the separated shoot, i.e., a segment size of perhaps 2 millimeters. The stem segment can then be fixed, and subjected to the GUS histochemical assay. The result is a cross-sectional view of the stem of the soybean plant which will exhibit a color, i.e., blue, in those portions of the tissues of the plant which have been transformed. Based on an analysis of those tissues, it is then possible to predict for those stems which exhibit some enzyme activity, those plants which are most likely to yield a germ line transformation event.

[0030] A classification scheme has been devised to categorize the results of the stem segment assay. The transgenic primary regenerates were designated "B" when the GUS enzyme assay revealed that the gene was expressing in all of the epidermis, the cortex, and the pith of the stem segments assayed. The regenerates were designated "C*" or "P*" when the expression of the GUS gene was confined to 100% of either the cortex or the pith respectively of the shoot segment which was assayed. The regenerating shoots were classified as "C" when at least half of the cortex

expressed the enzyme, and "c" when less than 50% of the cortex showed activity. Similarly a classification of P or p indicates GUS activity in more or less than 50% of the pith of the stem. The shoots were scored as "e" when the activity of the GUS gene could only be seen in the epidermis of the shoot. Examples were also observed in which the codes were used to indicate localized activity in multiple areas, for example, a score of c/P* indicates that the shoot expressed the GUS marker gene in 100% of its pith tissue, but in less than 50% of its cortex tissue. A classification of E*/p indicates the expression of the enzyme in 100% of the epidermis but in less than 50% of the pith and none of the cortex of the transformant plant.

[0031] Figures 3 - 8 illustrate the classification scheme of the stem segments. Fig. 3 is a highly stylized drawing of an ideal soybean stem cross-section in which 30 indicates the whole stem, 32 indicates the epidermis, which is greatly exaggerated in width, 34 indicates the cortex and 36 indicates the pith. Shading in Figs. 4 to 8 is intended to represent blue color in the GUS assay. Fig. 4 illustrates a stem 38 classified as "E" since only the epidermis is expressing the GUS gene. Fig. 5 illustrates a stem 40 which would be classified as "c/p" since less than 50% of cortex or pith are expressing. The stem 42 of Fig. 6 would be classified as "c/P*" since all the pith but less than 50% of the cortex was transformed. The stem 44 of Fig. 7 would be "C*". As should be apparent, the stem 46 of Fig. 8 represents a "B" stem segment. All of the results of Figs. 4 - 8 represent actual events.

[0032] For most of the shoots recovered for the plant transformation process using the apparatus of Figs. 1 and 2, no expression of the GUS gene in the stem section was found. Of those shoots in which expression in the stem segment was found, and classified in one of the categories as described above, a number of shoots of each category were either induced to root or were grafted onto healthy root stock and cultivated further into plantlets. When the plantlets developed either two or three trifoliate leaves, one or more leaflets from the first trifoliate leaf was harvested from each of the plants and the trifoliate leaflet itself was assayed for activity for the GUS enzyme. Thus new phenotypes could be identified which could be correlated to the phenotypes from the classification on the original stem of the primary regenerate shoot. A number of the plants were found only to express the GUS enzyme in epidermal trichomes and/or hairs either throughout the leaflet or only in certain areas within the leaflet. Other plants were found to express GUS activity only in mesophyll cells or stomata, and the activity in these tissues could extend throughout the leaflets or be confined to one or more specific areas within the leaflet. In other cases, the enzyme activity would be found to be localized in the mid-rib or to extend in varying degrees to one or both sides of the mid-rib in sectors of the leaflet. Examples were even found in which the GUS activity extended from the mid-rib to the perimeter only on one-half of the leaf. Again, the most optimal phenotype observed was that in which the blue color was not either localized into a sector or confined to the epidermis.

[0033] Upon further development, a third assay was developed based on petiole and mid-rib sections on a trifoliate leaf higher up the regenerating plant. It was found that if this last assay were not performed, certain non-germ line transformation events would be allowed to go to seed, since certain transformation events seem to give rise to plants in which the transformed tissues only extended a finite length up the stem of the regenerating plant. It became a not uncommon observation to observe GUS activity only in the first trifoliate leaf and not in certain other trifoliates further up the plant. It was, however, the general observation that the majority of plants which had GUS activity in the third or fourth trifoliate leaves would exhibit the same activity in all or most of the leaves of the plants.

[0034] The use of these three indicia led to the observation that the transformation process could be optimized to the total overall labor involved if only the plants which were characterized as "B", "P*" or some combination of P and c or C, during the assay of the stem segment, were fully regenerated and allowed to propagate into R1 plants. Even plants which are characterized as B, however, would not prove to have germ line transformation events if the expression of the GUS gene was not also found in the first trifoliate leaf and also in the petiole mid-rib assay in the mature leaf. The best use of the second and third indicia of transformation events were used as negatives, that is to say, plants which failed to properly exhibit the GUS activity at these stages could also be discarded, since the likelihood that such plants would yield germ line transformation events was found to be relatively minimal. Of plants which had been classified as B during the stem segment, and which showed major GUS activity in the pith or pith and cortex during the assay of the petiole and mid-rib sections of the third or fourth trifoliate stage, all plants had activity which was later determined to involve germ line transformation events as evidenced by the recovery of progeny plants which were clonal and expressing the GUS gene. All plants which were classified in any other of the categories described above exhibited a dramatically lower frequency of germ line transformation events, although such events did occur with other classes of tissues. For example, two plants out of 363 which showed activity only in their epidermis at the stem segment stage also gave rise to transformed progeny. However, the likelihood of any transformation events from any other categories of plants was thus so low as to justify, on a practical basis, discarding all but the plants designated above during the stem segment and further to sexually propagate those plants categorized likely to lead to germ line transformation which both exhibited GUS activity at the first trifoliate stage and also showed major pith and cortex involvement during the assay of the petiole and mid-rib of the mature leaf of the plant.

[0035] An excess of one-third of the plants classified as "B" would give rise to germ line transformation events. By contrast, plants categorized in other categories during the stem segment as they were found to have a dramatically

lower likelihood of germ line transformation event. For example, a plant that is classified as c/p*, the likelihood of a germ line transformation event was less than 10% while for plants showing activity in greater than half their cortex, and classified C, the ratio of transformation events was on the order of 2%. Of plants which had major cortex and pith activity, designated C/P, of which only eight plants were tested, two yielded germ line transformation events. Of six plants which showed activity in less than 50% of the pith, one germ line transformation event was recovered. Thus the exact classifications that are regenerated may depend on a cost analysis of the cost of regeneration of shoots into plants and seeds as compared to the cost of generating more putatively transformed shoots.

EXAMPLES

[0036] The target tissues used in the soybean transformation procedures described herein were the excised embryonic axes from immature and mature soybean seeds of elite lines of soybean. The axes were excised from the seeds and plated on target plates on agar. The methodology for excising the seeds and performing the transformation experiment is described in detail in McCabe et al., *Bio/Technology*, 6:923-926 (1988). The apparatus used in the transformation experiments was the apparatus of Figs. 1 and 2.

[0037] After the transformation event was conducted, in excess of 5,000 shoots were recovered, and subject to a stem segment assay. In performing the stem segment assay, the shoots were allowed to reach a length of approximately 2 centimeters before they were severed from the underlying tissue. At the time of severing, a 2 to 4 millimeter section of the base of the shoot was cut off, and assayed for GUS activity. Of the 757 stem segments assayed which showed GUS activity, the plants were categorized on the basis of the kind of activity which was observed in the stem by the categorization system described above, i.e., either B, C/P*, C, C/P, p, c or e. The distribution of the plants as classified can be seen in Table I.

TABLE I

Germ Line Event Frequency			
Stem Segment Assay Classification	Number of Plants	Number expressing in first leaf	Number expressing in mature leaf
B	30	27	13
C/P*	13	5	1
C	45	20	1
C/P	8	4	2
p	6	1	1
c	292	9	1
e	363	7	1

[0038] All of the plants for whom the stem segment assay was positive to any degree were then either grafted or rooted to create plantlets. The first trifoliate leaf from each of the plantlets was assayed for GUS activity. The results of that assay are indicated by the third column in Table I. All plants which continued to have GUS activity in the first trifoliate leaf were then propagated into plants containing multiple leaves, and mature third or fourth trifoliate level leaves were harvested from the plants and the petiole and mid-rib sections were taken through that leaf and were fixed and assayed for GUS activity. The results of those assays are indicated in the fourth column in Table I. All plants indicated in the fourth column were found to have experienced a germ line transformation event.

[0039] The fourth column in Table I indicates those plants which had activity in the pith or pith and cortex of the third or fourth stage trifoliate leaf. All of the plants which had GUS activity were allowed to self-pollinate and reproduce progeny. Analysis of the progeny was then done for GUS activity. It was determined that all of the plants which were indicated in column 4 of this chart to have activity in the pith or pith and cortex of the third or fourth trifoliate leaf stage were found to involve germ line transformation events as indicated by the recovery of progeny plants which expressed the GUS gene.

[0040] Of plants which failed to have pith or pith and cortex activity in the third or fourth stage trifoliate leaf, the results were more mixed. Of plants which had been designated B by the stem segment assay, during the assay of the third or fourth trifoliate leaf, 8 of the plants only showed activity in the epidermis while 6 were found to be completely negative in GUS assay. Two of the plants which showed only activity in their epidermis gave rise to transformed progeny.

- by the coding sequence in the cells of soybean plants;
- (b) coating copies of the exogenous genetic construction onto carrier particles of biologically inert material, which are small in size relative to the size of soybean cells;
- (c) accelerating the coated carrier particles into the cells of regenerable soybean tissue;
- (d) regenerating shoots from the soybean tissue;
- (e) severing a segment of the stem from each of the shoots;
- (f) assaying the stem segments for the expression of the gene product of the marker gene;
- (g) regenerating into whole sexually mature soybean plants only the shoots, the stem segments of which expressed the gene product of the marker gene in at least a major portion of its pith;
- (h) sexually propagating the whole plants to create progeny plants; and
- (i) assaying for the expression of the gene product of the marker gene in the progeny plants to indicate the germ line insertion of the gene of interest into the germ line of the progeny plants.
4. A method according to claim 3 wherein the marker gene is a gene coding for the enzyme beta-glucuronidase and where the assaying in steps (f) and (i) is done for the activity of that enzyme.
5. A method according to any one of the preceding claims wherein the carrier particles are gold particles.
6. A method according to any one of the preceding claims wherein in the accelerating step, the motive force is provided by an electric voltage spark discharge.
7. A method according to claim 6 wherein the accelerating step includes layering the carrier particles onto a carrier sheet, placing the carrier sheet near to a pair of spaced electrodes and then initiating an electric voltage spark discharge between the electrodes.
8. A method according to any one of claims 3 to 7 wherein the soybean tissue into which the carrier particles are accelerated comprises excised embryonic axes from soybean seeds.
9. A method according to claim 8 wherein the embryonic axes are plated after the particle accelerating step and in the regenerating step shoot formation is induced in the plated embryonic axes.
10. A method according to any one of claims 3 to 9 wherein each regenerating shoot is severed from the soybean tissue and at substantially the same time a stem segment is severed from each shoot.
11. A method according to any one of claims 3 to 10 wherein the regeneration step (g) is performed only for plants the stem segments of which expressed the marker gene in all the pith, cortex and epidermis, in more than 50% of the cortex and more than 50% of the pith, or in more than 50% of the pith, of the stem segment.
12. A method according to any one of claims 3 to 11 wherein the sexual propagation step (h) is performed by self-pollination.

Patentansprüche

1. Verfahren zur Keimbahntransformation von Bohnenpflanzen, wobei das Verfahren Schritte umfasst, bei denen man:
- a) ein exogenes genetisches Konstrukt erzeugt, das eine kodierende Sequenz und eine flankierende regulatorische Sequenz umfasst, die in der Lage ist, ein Genprodukt, das durch die kodierende Sequenz kodiert wird, in den Zellen der Pflanze zu exprimieren;
- (b) Trägerpartikel, die aus biologisch inertem Material bestehen und im Vergleich zur Größe der Pflanzenzellen klein sind, mit Kopien des exogenen genetischen Konstruktes beschichtet;
- (c) die beschichteten Trägerpartikel in die Zellen eines regenerierbaren Gewebes der Pflanze beschleunigt;
- (d) das Gewebe zu Pflanzen regeneriert;
- (e) während des Regenerationsverfahrens ausgewählte Teile des Stammes einer jeden regenerierten Pflanze

auf die Gegenwart des genetischen Konstrukts hin untersucht, wobei der ausgewählte Teil des Stammes einen Mark-Bereich umfasst und bei dem Nachweisverfahren bestimmt wird, ob das genetische Konstrukt in dem Mark-Bereich exprimiert wird;

- 5 (f) vorzugsweise solche Pflanzen sexuell vermehrt, deren ausgewählten Teile einen positiven Nachweis auf die Gegenwart des genetischen Konstruktes erbrachten, um Nachkommenpflanzen zu erzeugen; und
 - (g) die Gegenwart des genetischen Konstrukts in den Nachkommenpflanzen verifiziert.
- 10 2. Verfahren nach Anspruch 1, bei dem die Pflanzen Sojabohnen sind.
 3. Verfahren zur Erzeugung von Sojabohnenpflanzen, die in der Keimbahn transformiert sind, nach Anspruch 1, bei dem man:
- 15 (a) ein exogenes genetisches Konstrukt erzeugt, das für eine Insertion in Sojabohnen geeignet ist, wobei das Konstrukt Expressionskonstrukte sowohl für ein Marker-Gen als auch ein ausgewähltes Gen umfasst, wovon jedes eine kodierende Sequenz und eine flankierende regulatorische Sequenz umfasst, die in der Lage ist, das durch die kodierende Sequenz kodierte Genprodukt in den Zellen der Sojabohnen zu erzeugen;
 - 20 (b) Trägerpartikel aus biologisch inerten Material, welche im Verhältnis zur Größe der Sojabohnenzellen kleins sind, mit Kopien des exogenen genetischen Konstrukts beschichtet;
 - (c) die beschichteten Trägerpartikel in Zellen eines regenerierbaren Sojabohnengewebes beschleunigt;
 - 25 (d) Triebe aus dem Gewebe regeneriert;
 - (e) eine Segment des Stammes eines jeden Triebes abtrennt;
 - (f) das Stammsegment auf die Expression des Genproduktes des Marker-Gens hin untersucht,
 - 30 (g) nur solche Triebe, deren Stamm-Segmente das Genprodukt des Marker-Gens wenigstens in einem wesentlichen Teil des Markes des exprimieren, zu vollständigen, sexuell reifen Sojabohnenpflanzen regeneriert;
 - 35 (h) die ganzen Pflanzen sexuell fortpflanzt, um Nachkommenpflanzen zu erzeugen; und
 - (i) die Expression des Genproduktes des Marker-Gens in den Nachkommenpflanzen untersucht, um Keimbahninsertionen des gewünschten Gens in die Keimbahn der Nachkommenpflanzen zu identifizieren.
- 40 4. Verfahren nach Anspruch 3, bei dem das Marker-Gen ein für das Enzym β -Glukuronidase kodierendes Gen ist und bei dem das Verfahren in den Schritten (f) und (i) auf dem Nachweis der Aktivität dieses Enzyms beruht.
 5. Verfahren nach einem der vorstehenden Ansprüche, bei dem die Trägerpartikel Goldpartikel sind.
 - 45 6. Verfahren nach einem der vorstehenden Ansprüche, bei dem die antreibende Kraft in dem Beschleunigungsschritt durch Funkenentladung mittels elektrischen Stroms zur Verfügung gestellt wird.
 7. Verfahren nach Anspruch 6, bei dem der Schritt der Beschleunigung eine Ablagerung der Trägerpartikel auf einer Trägerplatte, die Platzierung der Trägerplatte nahe eines Paares von getrennten Elektroden und die Auslösung der Funkenentladung durch elektrischen Strom zwischen den Elektroden umfasst.
 - 50 8. Verfahren nach einem der Ansprüche 3 bis 7, bei dem das Sojabohnengewebe, in das die Trägerpartikel beschleunigt werden, freigelegte embryonale Achsen von Sojabohnen-Samen umfassen.
 - 55 9. Verfahren nach Anspruch 8, bei dem die embryonalen Achsen nach dem Partikel-Beschleunigungsschritt ausplattiert werden und bei dem während des Regenerationsschrittes die Triebbildung in den ausplattierten embryonalen Achsen induziert wird.

10. Verfahren nach einem der Ansprüche 3 bis 9, bei dem jeder der sich regenerierenden Triebe vom Sojabohnengewebe abgetrennt wird und ein Segment des Stammes im wesentlichen zum selben Zeitpunkt von jedem Trieb abgetrennt wird.
- 5 11. Verfahren nach einem der Ansprüche 3 bis 10, bei dem der Regenerationsschritt (g) nur mit solchen Pflanzen durchgeführt wird, deren Stammsegmente das Marker-Gen sowohl in Mark, Rinde und Epidermis exprimieren, und zwar in mehr als 50 % der Rinde und mehr als 50 % des Markes oder in mehr als 50 % des Markes des Stammsegmentes.
- 10 12. Verfahren nach einem der Ansprüche 3 bis 11, wobei der sexuelle Fortpflanzungsschritt (h) durch Selbstbestäubung durchgeführt wird.

Revendications

- 15 1. Méthode pour la transformation de la lignée germinale de plants de haricots, laquelle méthode comprend les étapes de :
- 20 (a) préparer une construction génétique exogène comprenant une séquence codante et une séquence régulatrice adjacente efficaces pour exprimer un gène produit codé par la séquence codante dans les cellules de la plante ;
- (b) revêtir des copies de la construction génétique exogène sur des particules de support qui sont en un matériau biologiquement inerte et qui sont de petites dimensions relativement aux dimensions de cellules de la plante ;
- 25 (c) accélérer les particules revêtues de support dans les cellules de tissu régénérable de la plante ;
- (d) régénérer le tissu dans les plantes ;
- (e) pendant le procédé de régénération, analyser une portion sélectionnée de la tige de chaque plante en régénération pour la présence de la construction génétique, la portion de tige sélectionnée comprenant une section médullaire, l'analyse déterminant si la construction génétique est exprimée dans la section médullaire ;
- 30 (f) multiplier sexuellement préférentiellement les plantes dont les portions sélectionnées sont positives pour la présence de la construction génétique pour créer des plantes en descendance ; et
- (g) vérifier la présence de la construction génétique dans les plantes en descendance.
2. Méthode selon la revendication 1 où les plantes sont des plantes de soja.
- 35 3. Méthode selon la revendication 1 de création de plantes de soja à lignée germinale transformée comprenant les étapes de :
- 40 (a) préparer une construction génétique exogène appropriée pour une insertion dans des plantes de soja, laquelle construction comprend des constructions d'expression pour un gène marqueur et un gène d'intérêt, dont chacun comprend une séquence de codage et une séquence régulatrice adjacente efficace dans des cellules de soja pour produire un gène codé par la séquence de codage dans les cellules des plantes de soja ;
- (b) revêtir des copies de la construction génétique exogène sur des particules de support en un matériau biologiquement inerte, qui sont de petites dimensions relativement à la dimension des cellules de soja ;
- 45 (c) accélérer les particules de support revêtues dans les cellules du tissu de soja régénérable ;
- (d) régénérer des pousses à partir du tissu de soja ;
- (e) rompre un segment de la tige de chacune des pousses ;
- (f) analyser les segments de tige pour l'expression du gène produit du gène marqueur ;
- (g) régénérer dans des plants entiers sexuellement murs de soja uniquement les pousses, les segments de tige dont est exprimé le gène produit du gène marqueur dans au moins une portion majeure de sa moelle ;
- 50 (h) propager sexuellement les plants entiers pour créer des plants en descendance et
- (i) analyser l'expression du gène produit du gène marqueur dans les plants à descendance pour indiquer l'insertion de la lignée germinale du gène d'intérêt dans la lignée germinale des plantes en descendance.
4. Méthode selon la revendication 3, où le gène marqueur est un gène codant pour l'enzyme bêta-glucuronidase et où l'analyse des étapes (f) et (i) est faite pour l'activité de cette enzyme.
- 55 5. Méthode selon l'une quelconque des revendications précédentes où les particules de support sont des particules d'or.

6. Méthode selon l'une quelconque des revendications précédentes où dans l'étape d'accélération, la force motrice est produite par une décharge d'étincelle de tension électrique.
- 5 7. Méthode selon la revendication 6, où l'étape d'accélération comprend la mise en couche des particules de support sur une feuille de support, la mise en place de la feuille de support près d'une paire d'électrodes espacées puis l'initiation d'une décharge d'étincelle de tension électrique entre les électrodes.
- 10 8. Méthode selon l'une quelconque des revendications 3 à 7, où le tissu de soja dans lequel les particules de support sont accélérées comprend des axes embryonnaires excisés des graines de soja.
- 15 9. Méthode selon la revendication 8, où les axes embryonnaires sont plaqués après l'étape d'accélération des particules et dans l'étape de régénération, la formation des pousses est induite dans les axes embryonnaires plaqués.
- 20 10. Méthode selon l'une quelconque des revendications 3 à 9 où chaque pousse régénérante est rompue du tissu de soja et sensiblement en même temps un segment de tige est rompu de chaque pousse.
- 25 11. Méthode selon l'une quelconque des revendications 3 à 10, où l'étape de régénération (g) est accomplie uniquement pour les plantes dont les segments de tige expriment le gène marqueur dans la moelle, le cortex et l'épiderme, à plus de 50% du cortex et plus de 50% de la moelle ou à plus de 50% de la moelle, du segment de tige.
- 30 12. Méthode selon l'une quelconque des revendications 3 à 11, où l'étape (h) de propagation sexuelle est accomplie par auto-pollinisation.
- 35
- 40
- 45
- 50
- 55




FIG. 1





FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8